

Curry County Agricultural Water Quality Management Area Plan

Developed by

The Oregon Department of Agriculture

and

the Curry Soil and Water Conservation District

with assistance from

the Curry County Local Advisory Committee

June 3, 2004

Local Advisory Committee Members

**Walt Schroeder, Chair
Knute Andersson
Joe Brown
Becky Crockett
Jim Donaldson**

**George Fleming
Frank Burris
Harry Harms
Mike Knapp
Earl Lang**

**Lee Riddle
Jim Waldvogel
Rick McKenzie
Robert McKenzie
Ted Fitzgerald**

Blank

Table of Contents

<i>An Open Letter from the LAC to the Curry Agricultural Community</i>	1
<i>Foreword and Applicability</i>	2
<i>Acronyms</i>	3
<i>History of 303(d) list and Senate Bill 1010</i>	4
<i>Geographic and Programmatic Scope</i>	5
<i>Geographic Uniqueness within Curry County</i>	5
Agriculture.....	5
Fisheries.....	7
The Watershed as an Ecosystem.....	7
Climate.....	8
<i>Physical Settings of Curry County Watersheds (north to south)</i>	9
Floras Creek.....	9
Sixes.....	10
Elk River.....	12
Hubbard Creek.....	13
Euchre Creek.....	13
Rogue River.....	14
Hunter Creek.....	15
Pistol River.....	16
Chetco River.....	17
Winchuck River.....	18
<i>Mission</i>	20
Goal, Objective, Responsibility, and Intent.....	20
Goal of the Committee:.....	20
Objectives:.....	20
Responsibility of the Committee as stated in ODA Policy:.....	20
<i>Water Quality Issues of Concern</i>	21
Introduction.....	21
Contributing Factors.....	21
303(d) list -Addressing 303(d) listed Parameters.....	21
The Coastal Zone Act.....	22
<i>Prevention and Control Measures</i>	23
<i>Strategies for Implementation</i>	27
<i>Monitoring and Evaluation</i>	27
Baseline Condition Monitoring.....	28
Water Quality Trend Monitoring.....	28
Effectiveness Monitoring.....	28
Site-Specific Rule Compliance Monitoring and Assessment.....	28
Current and future monitoring and assessment efforts.....	28
<i>Education and Outreach</i>	29
<i>Administrative Roles and Responsibilities</i>	31
Total Maximum Daily Loads.....	31
Designated Management Agency/Local Management Agency.....	31
Monitoring and Evaluation of the Plan’s Effectiveness.....	31
<i>Complaints and Inspections</i>	32
Compliance Procedures.....	32
Review Period.....	32
<i>References</i>	35
<i>Appendix A. Available Technical and Financial Assistance</i>	37
<i>Appendix B. Coastal Zone Management Act Measures</i>	38
<i>Appendix C. Enforcement Procedures</i>	40
<i>Appendix D. Map of Curry County Agricultural Water Quality Management Area</i>	43

An Open Letter from the LAC to the Curry Agricultural Community

The Curry Local Advisory Committee (LAC), after discussing the riparian rule at every meeting since February 2002, developed the following riparian statement at the March 12, 2003, meeting.

Riparian Management

A Riparian area is an edge or bank of a river, tributary, or other body of water.

A proper functioning riparian condition is one important component of clean water and healthy watersheds.

Agricultural management complements the proper functioning condition of a riparian area by allowing vegetation and biomass accumulation that benefits water quality.

Upon rule adoption, management activities in the riparian area of anadromous fish bearing streams, that are 303d listed for temperature, will be conducted in a manner that avoids increases in stream temperature.
Exemptions shall include stream crossings, access for irrigation equipment and other accepted water dependent agricultural uses when conducted in a manner that minimizes impacts on streambank stability.

The Oregon Department of Agriculture (ODA) rejected this suggestion as inadequate and vulnerable and has inserted its own wording for a riparian rule. A majority of the Curry LAC believes this rule to be adequate to protect watersheds and water quality, while at the same time protecting private property rights.

Sincerely,

Walt Schroeder, Chair, Curry LAC

Foreword and Applicability

This Agricultural Water Quality Management Area Plan (Area Plan) provides guidance for addressing agricultural water quality issues in the Curry County Agricultural Water Quality Management Area (Management Area). The purpose of this Area Plan is to identify strategies to reduce water pollution from agricultural lands through a combination of educational programs, suggested land treatments, management activities, and monitoring. The provisions of this Area Plan do not, by themselves, establish legal requirements or prohibitions. The Oregon Department of Agriculture will exercise its enforcement authority for the prevention and control of water pollution from agricultural activities under administrative rules for the Curry County Area Plan and Oregon Administrative Rules (OAR) 603-090-0120 through 603-090-0180.

The administrative rules for the Curry County Basins set forth the requirements and/or prohibitions that will be used by ODA in exercising its enforcement authority for the prevention and control of water pollution from agricultural activities. In addition, OARs 603-090-0120 through 603-090-0180 describe the enforcement actions that may be triggered upon the finding of a violation by ODA.

Acronyms

AgWQM - Agricultural Water Quality Management

Area Plan - Curry County Agricultural Water Quality Management Area Plan

Area Rules - Curry County Agricultural Water Quality Management Area Rules

DEQ - Oregon Department of Environmental Quality

LAC - Local Advisory Committee

NRCS - Natural Resource Conservation Service

OAR - Oregon Administrative Rule

ODA - Oregon Department of Agriculture

ORS - Oregon Revised Statutes

OSU - Oregon State University

OWT - Oregon Water Trust

SB 1010 - Senate Bill 1010

SWCD - Soil and Water Conservation District

TMDL - Total Maximum Daily Load

VOC - Volatile Organic Chemical

History of 303(d) list and Senate Bill 1010

In 1972, President Nixon signed into law the Clean Water Act. This was a time in our history when rivers were catching on fire (the Cuyahoga in Ohio), large lakes were completely sterilized (Lake Erie), and there were massive estuarine die-offs (Chesapeake Bay). Point source discharges from factories and treatment plants were targeted for clean up and Total Maximum Daily Loads (TMDLs) were established for contaminants that threatened the well being of our population and our natural resources. In 1999, Lake Erie and the Chesapeake Bay have a thriving fishery while the Cuyahoga River has people swimming and boating its waters. The problem is that there are still large amounts of pollutants entering waterways with no discrete source. This type of pollution is called non-point source pollution.

In Oregon these non-point sources were the focal point of a suit filed in 1986 charging that Oregon was not enforcing the Clean Water Act standards thus harming both fish and human populations. The court agreed and charged the Department of Environmental Quality (DEQ) with cataloging the waters of the state that were impaired as characterized by the beneficial uses they supported. The list became known as the 303(d) list (available at <http://www.deq.state.or.us> or local DEQ office 541-269-2721).

In an effort to insure that the agricultural industry's unique concerns and constraints were addressed under state and federal water quality drivers, industry representatives lobbied the state legislature for a possible solution. The 1993 Oregon Legislature, in passing Senate Bill (SB) 1010, provided for the Oregon Department of Agriculture to be the lead state agency to address water pollution coming from private agriculture land. Under the law, the ODA is authorized to develop and carry out a water quality management plan for any agricultural or rural lands where a water quality management plan and its associated rules are required by state or federal law. TMDL development in a basin encompasses all the potential nonpoint pollution sources, of which agricultural is but one part.

Geographic and Programmatic Scope

The South Coast Agricultural Water Quality Management Area Plan (Area Plan) boundaries include the California border to the south, Josephine County boundary to the east, the Pacific Ocean on the west, and the southern watershed boundaries of the Coos/Coquille Management Area on the north. The northern boundary of the South Coast plan includes parts of the county line but includes all of the Floras Creek drainage and none of the Two Mile and Four Mile Creek drainages.

In 1995, the Oregon legislature recognized potential conflicting authorities that belonged to both ODA and DEQ. While SB 1010 gave the recognition of water quality problems to ODA, DEQ still had authority to regulate any water quality problem with its authority under Oregon Revised Statute (ORS) 468B if an Agricultural Water Quality Area Plan did not achieve standards adopted by the Environmental Quality Commission. SB 502 in 1995 clarified that conflict and firmly authorized ODA to conduct all water quality regulation that occurred on agricultural lands for agricultural practices. A 1996 opinion from the Oregon Attorney General's office states that ODA has the statutory responsibility to regulate agriculturally related pollution. That same opinion also recognized the need to define that authority by developing water quality plans and rules that specifically addressed agricultural practices and land conditions and achieve the standards adopted by the Environmental Quality Commission.

ORS 468B.010 to 468B.050 lays out a broad framework under which pollution is defined as both point and non-point sources which degrade water quality so as to be detrimental to beneficial uses. State water quality standards are based on those beneficial uses (i.e. 64 degree temperature standard, 406 colonies of E. coli bacterial standard). The broad nature of the statute places a heavy burden of responsibility and confusion on the private agricultural landowner. How was the landowner going to determine if his/her operations were harmful to the beneficial uses of a particular water body? Area Plan and Rule-making seeks to eliminate the confusion of landowners and identify the responsibility of the agriculture community by working with Local Advisory Committees (LAC) to develop land conditions that describe to both ODA and the landowner what the land looks like that has a potential to cause a 468B violation. In this scenario, ODA retains its full authority to interact with agricultural landowners for resource protection but landowners are the ones who identify the unique conditions that can exist in a particular watershed thereby protecting their operations from unnecessary scrutiny even if local waters were in violation of water quality standards.

Geographic Uniqueness within Curry County

Agriculture

The first evidence of agriculture in Curry County was in 1851 when Captain William Tichenor, the founder of Port Orford, brought six horses and some swine and established a settlement at that place. The livestock and associated crops, as well as gardens to provide vegetables for the increasing number of settlers, increased rapidly.

The rich virgin soils in Curry County and year around grazing and mild climate were ideal for raising livestock and some crops. By 1880 the assessment roles showed 22,000 sheep and close to 4000 cattle grazing the lush mountain prairies and river bottomlands. Close to 800 hogs ranged through the forests and mountain prairies feeding on roots and mast. (Tanoak acorns and Myrtle nuts). Lightning fires had burned much of the area. Some had been burned by the Native Americans to provide habitat for deer and elk. These open mountain prairies were ideal for sheep and cattle raising.

The river bottom lands and associated hill-grazing lands made dairy farming the number one farm enterprise in the county. Cheese factories in almost every community manufactured cheese and butter to ship to San Francisco and other markets. A nationally famous Blue Cheese factory was located in Langlois until destroyed by fire in 1957. Close to 400 dairy herds were extant in the county in 1938, though at that time many were small herds that could be hand milked by the farmer and his children. Now there are only two dairy herds in the entire county. Both have waste management plans and are regulated under the ODA Confined Animal Feeding Operation program (CAFO).

The early sheep were raised primarily for their wool. Curry wool was prized for its consistent growth and long fiber. In the 1930s Curry listed 25,000 head of mature sheep. But soon, with cross breeding and better management, the market lamb industry provided additional income to sheep producers. In recent years predators and low prices for lamb and wool, have closed down many sheep operations. Several sheep operations are now devoted only to cattle.

During World War II, the Easter lily enterprise greatly expanded when bulbs were no longer available from Japan. Hundreds of farmers and gardeners got into the lily business. The industry started in the Bandon area but over the years it centered on the rich deep soils and climate of the Harbor Bench and nearby Smith River delta in northern California. Growers have developed ponds or wells from which they irrigate their lilies. Easter lilies are a crop unique to the soils and climate of this area and are a stable industry, providing many jobs.

Cranberries were introduced to the south coast in 1885 at Hauser in Coos County and expanded to south Coos County in the Bandon area. Soon the most rapid cranberry acreage being developed was in north Curry County. In 1996 Curry County accounted for 25 percent of the production in Oregon. Cranberry production uses water for frost protection, irrigation, weed and pest control, temperature control, and for harvest. The preferred harvest method is to flood the beds and beat the vines to separate the berries from the vines. The berries float and are corralled for loading onto trucks. Dry harvest is also used but it is not a preferred method. Cranberry growers possess water rights to apply water and most have constructed reservoirs to hold surplus winter rainfall. A majority of growers recycle water through a series of beds at slightly different elevations, reducing the use of water from springs and creeks.

In 2000 the cranberry industry suffered a severe setback when stocks of cranberries in storage were at an all time high. The cost of production was considerably more than the price received by the growers.

Several other crops are grown on the rich soils of the county including hydrangeas, rhododendrons and blueberries. The largest hydrangea farm in the world is located on the Harbor Bench in Southern Curry County.

The agricultural history of northern Curry County is unique. Small streams in the lower portion of the streams were channelized and straightened from the mid 1920s through the 1950s. This work was done through various federal programs (such as Swampbuster) and by private landowners. The reason for stream alteration was to capitalize on the rich bottomlands that supported the many large dairies and livestock operations in North Curry. The affected streams include Langlois Creek, Morton Creek, Bethel Creek, Croft Lake outlet, New Lake outlet (Bono Ditch), and portions of Willow Creek. The lower portions of these streams function as agriculture drains. Historically, they have been cleaned and maintained as needed by the individual landowners.

Approximately two thirds of the county is publicly owned: U.S. Forest Service, Bureau of Land Management, State of Oregon, and County. With limited privately owned land in the county, pressure from development is causing concern as homes and other structures are impinging on farmland.

Fisheries

Except for the streams with impassable barriers (such as the Harbor Bench area), all Curry County streams (from Floras Creek in the north to the Winchuck River in the south) are excellent producers of salmonids: Chinook salmon, steelhead, cutthroat trout, and Coho salmon. Local watershed councils, the Curry County Soil and Water Conservation District, and the Salmon Trout Enhancement Program are working to enhance most of these streams and address impacts created by past land use practices. Local agricultural producers have worked with these groups to enhance riparian and upland areas and proactively improve watershed health.

Changes in some native salmonid populations have occurred in Curry County rivers but not to the extent experienced in counties to the north and south. Factors influencing the changes vary between species and river systems, but include spawning and rearing habitat degradation, passage obstacles, overfishing, water demands for other uses, urban development, seasonally limited streamflow and probably most important, changing ocean habitat conditions (El Niños, upswelling, ocean oscillations). However, we should not judge yesterday's activities by today's standards, but only recognize that everyone has contributed to the declines and move forward to correct past mistakes.

The nine watershed councils in Curry County have contributed greatly to the enhancement of salmonid habitat through riparian fencing, planting, improving fish passage, instream structure development, habitat improvement, etc. Ranchers have fenced many stream miles voluntarily under the Hire the Fisher program and the Oregon Plan for Salmon and Watersheds. As a result of these efforts, as well as improvements in natural conditions, the abundance of several salmonid species in Curry County streams is the highest in Oregon and in some cases close to historic levels.

Salmonids adapted in ecosystems that historically had a high degree of stream complexity including large woody debris, flood plains, good spawning gravel reserves, estuaries, and wetland refugia, meandering stream courses and in some cases lake systems in the watershed. Human activities had altered some of this traditional salmonid freshwater habitat.

El Nino events, relatively dry winters for 4-5 years and the past 20 years of altering ocean conditions have had an adverse effect on anadromous fish populations. In recent years favorable ocean conditions and wet winter cycles have occurred and we are seeing quantum increases in salmonid returns to many Curry County streams. This suggests that ocean productivity is a long-term natural cycle that falls outside the realm of controllable fisheries management. Some declines in numbers of native fish have conflicted with increases in hatchery-raised stocks. Targeting hatchery fish while allowing native stocks to spawn naturally in the improved habitat may help to alleviate some of these concerns.

During periods when ocean productivity is low it is critically important to protect and improve freshwater habitats so they do not become the "weak link" in the chain of salmonid production. Agricultural operators actions can have an important impact on salmonid production and its freshwater habitat and these actions will be discussed in detail in this plan.

The Watershed as an Ecosystem

An ecosystem is an interdependent community of living and non-living organisms/elements including humans. Ecosystems do not always have defined boundaries. An ecosystem is a natural system composed of living and non-living elements working together to maintain conditions that support life.

A watershed is any area of land that drains water to a specific point such as a lake, river, or ocean. Like ecosystems, watersheds may be as large as the basin of the Mississippi River or as small as the water that flows into a pond. All land is in a watershed. In the hydrologic cycle, precipitation falls everywhere and drains somewhere. Energy inputs of sunlight, wind, and the hydrologic cycle interact with the landforms and the living species in ways that affect both the quality and quantity of water.

In an ideal condition, water is captured by infiltration into the spongy layer of duff and topsoil in the watershed. Some is held and by capillary *action* is available to plants. The remainder percolates down through the soil profile to recharge ground water supplies. The primary watershed process is the capture, storage, and slow release of water. This process helps to prevent excessive flooding during heavy winter rainfall and provides water in times of low rainfall. Where there is no topsoil, or where topsoil has been compacted, eroded, covered over by asphalt, buildings or concrete, or over-saturated, water is not captured but is allowed to run off over the surface of the ground and into watercourses. Flooding and turbidity is increased and water may not be available when needed during drier times of the year. The quality of water is improved by passage through the topsoil, which acts as a filter and serves as storage to increase base flow in summer.

Different landscape types within the watershed have different roles in the capture, storage, and slow release of water. Wetlands and flood plains slow down the movement of water allowing time for groundwater recharge. Wetlands also serve the purpose of filtering out possible contaminants. Vegetation helps hold topsoil in place and is an important source of humus in the form of decaying plant material. Healthy topsoil is not only our source of food; it also helps provide clean and abundant water.

Major fires historically occurred in the Klamath mountains eco-region on about 50-year intervals. Native Americans and ranchers both used fires to maintain grasslands and prairies. From the 1920s to 1987, the United States Forest Service adopted a policy to aggressively fight forest fires. Although no official policy change had taken place prior to 2002, current forest service practice is to allow fires to burn until they threaten private homes and structures. This first came to light with the Silver Fire in 1987, which burned thousands of acres in Curry County. In 2002, the Biscuit Fire confirmed this new practice as nearly 500,000 acres were consumed. Both national forest and private forest holdings were destroyed in that fire. Sufficient resources to stop this fire were not employed until substantial private residences and structures were put at risk. This watershed management practice is mentioned in the document only because the LAC feels that it has direct impact on stream temperature and sedimentation. We also feel that it has potential for substantially more long-range impact on fish populations in our watersheds.

Climate

Temperatures in Curry County are mild throughout the year because of the moderating influence of the Pacific Ocean. Along the coast, the difference between January and August average temperatures is only 14 degrees.

Annual rainfall averages vary from 75 or 80 inches along the coast to more than 100 inches on the upper slopes of the Coast Range. More than 70 percent of the annual precipitation falls between November and March, with an average of only 3 inches of rain falling during June, July, and August. During the winter, storms can produce intense rainfall; at least once per year, nearly four inches of rainfall can be expected during a 24-hour period.

Along the coast, there is less than 1 inch per year average snowfall. Inland, as many as 12 inches of snow have been recorded in a year, but there is frequently little to no snow accumulation. Snow usually melts within a few hours or days.

Strong winds occur frequently in Curry County. Peak gusts between 60 and 100 miles per hour can be expected a few times every year. During the summer, strong easterly winds bring warm, dry air to parts of the county, depleting soil moisture and often creating severe fire danger (Buzzard and Bowsby, 1970).

Physical Settings of Curry County Watersheds (north to south)

Floras Creek

The Floras Creek Watershed is an 83 square mile area located at the north edge of Curry County. Floras Creek crosses Highway 101 immediately south of the community of Langlois. Floras Creek provides *the* public water supply at Langlois as well as stock and irrigation water for the agricultural activities in the lower portion of the watershed. The upper portion of the watershed, most of the area east of Highway 101, is devoted to timber production and livestock grazing. Agricultural activities in the watershed area include livestock grazing, a dairy operation, and cranberry production. Over ninety percent of the land in the watershed is privately managed. The Bureau of Land Management manages five percent and the State of Oregon manages one percent.

The Floras Creek system includes five tributaries and Floras Lake. Boulder Creek, Willow Creek, Joe Cox Creek, Dwyer Creek, and White Rock Creek feed the north, east, and main forks of Floras Creek. Rainfall ranges from 70 to 80 inches along the coast and 90 to 115 inches at the higher elevations. Most of the rainfall comes as short intense storms from November through March. Less than ten percent of the rainfall occurs during the summer. These rainfall patterns, relatively small watershed, and limited water storage capacity result in large fluctuations in stream flow.

Extreme flooding is a defining feature of the watershed with floods occurring on a regular basis. High intensity rainfall events caused major floods in 1955, 1964, and 1983 and again in late 1996. The mouth of Floras Creek fluctuates in location almost annually. Floras Creek hits a foredune and flows north from 0 to 9 miles before breaching the foredune and exiting to the ocean. The fluctuating outlet location and the high flow events have contributed to severe separation of the stream (down-cutting) from its riparian area in the lower portion of the watershed. This separation and high flow events have contributed to streambank erosion, the cutting of wide stream cross-sections, and excessive deposition of gravel bars. The stream fluctuates between a meandering gravel-sand bed stream (Rosgen C4-5) to a straight entrenched stream (Rosgen F4-5) (Rosgen, 1996). During the dry summer months, these same highly eroded stream channels result in a relatively small stream flowing in a large channel. Thus, wide, shallow conditions contribute to low velocities and severe stream warming.

A large portion of the Floras Creek watershed is managed for timber and regulated under the Oregon Forest Practices Act. Most of this land was logged within the past forty years and consists of younger timber stands. Douglas fir is the most common tree used in reforestation and plantations. Some of these areas were left to naturally regenerate resulting in alder dominated stands. Some of the upland areas are managed for livestock grazing and these practices encourage vegetated cover throughout the year. There are also a few rural home sites.

Soils in the uplands are naturally steep, highly erodible, and prone to landslides. These natural geologic conditions in combination with historic logging practices and road construction (prior to Forest Practices Act), and various historic State, County, and private land and road management have contributed to an overall sediment load in the system. Present forest practices and other land and road management minimize upland erosion.

The lower portion of the watershed tends to be coarse to medium textured soils that are level and used for agricultural production. Extensive areas of these soils are allocated to improved pastures for ranching. Cranberries and some row crops are also grown. There is one dairy on Floras Creek. When the dairy is in use, the cows graze the pastures and are brought to the barn for milking. The highly erodible alluvial material presents streambank erosion problems for local landowners. Conservation minded landowners attempt to control loss of pasture and agricultural land by placing riprap, rock groins, bank barbs, or spruce trees along the banks. Unfortunately, these steps sometimes prove inadequate and the control structure is removed by floodwaters at times of severe flooding and down cutting. The landowners, in cooperation with the Bureau of Land Management, have submitted permits to mechanically breach the foredune in three locations to create a more consistent outlet for Floras Creek. A more consistent outlet and good vegetative conditions will help stabilize lower Floras Creek and prevent excessive erosion.

Some cranberry production exists in the Floras Lake sub-watershed and lower mainstem of Floras Creek. Cranberry production requires irrigation in the summer and flooding at harvest time in October. Both groundwater and surface water are used. The producers build storage ponds and recycle as much water as possible. The Oregon Water Resources Department and cranberry organizations in the area are cooperating on a water use study analyzing the relationship between ground water and surface water in the area. This study will help determine the potential impacts of groundwater use in cranberry production on low flow stream conditions.

Floras Creek and its tributaries are home to four species of anadromous fish and several species of freshwater stream, estuarine and marine fish species. Most of the species are native to the region, but some; large mouth bass and rainbow trout were introduced to enhance recreational fishing in Floras Lake. Coho salmon, Fall Chinook salmon, Winter Steelhead, and Cutthroat trout have historically spawned in the Floras Creek watershed (Maguire, 2001a).

Sixes

The Sixes River is one of the larger watersheds in the Southern Oregon coastal area, draining approximately 85,650 acres or 134 mi². Elevations in the watershed range from sea level to 3,315 ft. Major tributaries include the North, Middle and South Forks, and Dry, Edson, and Crystal creeks. The upper portion of the basin is characterized by steeply sloped forested land with narrow valleys and tributary streams that have moderately steep to very steep gradients. The predominant land use in the middle and upper portions of the watershed is commercial timber production. The lower few miles of the river are relatively low gradient coastal floodplain. Rural residential development, grazing, and other agricultural uses are the dominant land uses in the lower basin.

Approximately 92% of the Sixes River watershed is located in the Southern Oregon Coastal Mountains ecoregion, which is characterized by complex geology; steep, high gradient streams; and seasonally abundant precipitation. Watersheds in this ecoregion have high stream densities due to heavy precipitation during winter months. Extensive erosion can result from fractured geology, extremely variable stream flows, and a naturally high incidence of landslides in this ecoregion. The remaining 8% of the Sixes River drainage is located in the Coastal Lowlands ecoregion and is characterized by low gradient streams that are predominantly underlain by marine terrace deposits and flow through deep soils that range from silty clay loams to sandy loams. Streams in the Coastal Lowlands ecoregion are also susceptible to high erosion rates due to extreme fluctuations between summer and winter stream flows, easily eroded soils, incised channels and historic loss of riparian vegetation.

The predominant vegetation types found in the Sixes River watershed are listed in the table below. Natural disturbances that are capable of removing vegetation range from relatively frequent high winds and floods, to relatively infrequent forest fires, and even less frequent earthquakes. Potential riparian vegetation on low gradient lands may include dense thickets of wind-stunted shore pine,

Sitka spruce, and other brush species. Beaver are commonly found in low gradient channels of the Sixes River, and may significantly alter both vegetation and channel morphology.

Conifers	Sitka spruce, shore pine, grand fir, Douglas fir, western hemlock, Port Orford cedar and Monterey cypress
Hardwoods	Red alder, big-leaf maple, myrtle, and madrone
Shrubs	Rhododendron, holly, wax myrtle, willows, and Ceanothus spp.
Understory	Azalea, Ribes spp, iris, sea watch, huckleberry, salal, ferns, skunk cabbage, rushes, sedges, and grasses
Noxious	Gorse, Himalayan blackberry, tansy, scotch broom, European beach grass, and thistles

The Sixes River watershed has the greatest diversity of channel habitat types and more miles of low and moderate gradient channels than any watershed surveyed in Curry County. However, 41% of the watershed is composed of steep to moderately steep gradient channels. The risk ratings for increased erosion from roads within the Sixes River watershed range from low to high, with Dry Creek having the highest proportion (100%) of roads on slopes greater than 50% of any drainage on the South Coast of Oregon.

Approximately 69% of the land in the Sixes River basin is in private ownership, with 42% of the land owned and managed by private industrial timber companies and 27% owned by small-acreage, private landowners. Public lands comprise almost 29% of the lands in the basin, with management responsibility for these lands falling mostly on the U.S. Forest Service. Less than 3% of the land in the Sixes River watershed is in state, city, or county ownership.

Forestry is the dominant land use in the Sixes River watershed, accounting for 93% of the total land area. The remaining 7% of the watershed is used for agriculture, animal range, and rural residential development. Rangelands are managed for livestock grazing, with cows and sheep comprising the majority of livestock grazed.

Riparian habitats in Sixes River are characterized by a scarcity of large conifers near the surveyed channels and a relatively low incidence of bank erosion. Aquatic habitat complexity is low, primarily because of the lack of secondary channels and large wood throughout the drainage. Sixes River is listed as water quality limited by the Oregon Department of Environmental Quality from the mouth to the headwaters (including the South Fork Sixes) due to summer temperatures that exceed the 64^o F. temperature standard. Results from quarterly water sampling taken upriver from predominant agricultural lands on the Sixes River by DEQ indicate that water quality is moderately impaired due to high levels of nitrates, phosphates, and fecal coliform bacteria.

Out-of-Stream water rights for the Sixes River watershed consist of both storage rights (approximately 426 acre feet) and in-stream rights (120 cubic feet per second [cfs]), and the majority of those rights are senior to the 1964 In-Stream water right of 30 cfs in July and 25 cfs in August and September. Out-of-Stream water rights currently exceed the flow in Sixes River from May to October, with the water rights allocated for mining (75 cfs), agriculture (23 cfs), and irrigation (21 cfs) use. Water Storage Rights within the Sixes River watershed total 426 acre-feet and are entirely allocated for agricultural and irrigation use. Many of these rights are not exercised resulting in year-round flow.

An estimated 1,373 acres of wetlands have been assessed in the Sixes River drainage. Most of the wetlands in the drainage are found in the lower gradient sections along the lower mainstem of Sixes River. Approximately 124 of these acres have been highly modified, 837 acres have been

moderately modified, and the remaining 412 acres are not significantly altered. Wetlands within the Sixes River watershed are bordered primarily by forest and agricultural land, with a small proportion bordering rural residential, and other developed lands.

Chinook and Coho salmon, steelhead, and cutthroat trout are all native to the Sixes River drainage. The historic abundance and distribution of these salmonids within the watershed are not known. Chinook salmon and steelhead are thought to occupy most of their historic habitat in the Sixes River watershed. Coho salmon are listed coast-wide as threatened under the Endangered Species Act. Little is known about the historic distribution and abundance of cutthroat trout, but currently they are thought to fully utilize the available habitat. There are no hatcheries on the Sixes River yet the Sixes River continues to provide a thriving recreational fishery (Maguire, 2001b).

Elk River

The Elk River Watershed encompasses 59,520 total acres. The area has a maritime climate with annual precipitation from 90 inches at Anvil Creek to 130 inches in the headwaters. Approximately 80 percent of the precipitation occurs from October to March, and four percent during June, July and August. A small portion of the watershed lies between 2,400 and 4,600 feet in elevation, within a transient snow zone.

The River leaves the Siskiyou National Forest through a broad valley, and enters the ocean through a small estuary. Coastal lowlands make up 11 percent of the basin. Gradients are fairly low, fog and strong winds are common, and rainfall averages 60-90 inches per year. The average annual water yield is estimated to be 267,000 acre-feet. Low mean monthly flows of 20-100 cubic feet per second (cfs) occur between June and October, and high flows of 1,000-6,000 cfs occur between November and April. Peak flows of a magnitude greater than the 10-year return interval occurred in 1944, 1955, 1964, 1971, 1982, and 1995-96. The December 1964 flow was estimated to have an 80-150 year interval.

The Elk River watershed is comprised of Rocky Point sandstones and siltstones, Humbug Mountain conglomerate, shales of the Galice Formation, diorite intrusions, and ultramafic rocks. Vegetation in the watershed includes forested land with a hardwood/conifer mixture of Douglas fir, western hemlock, Port Orford cedar, Jeffrey pine, Sitka spruce, tanoak, red alder, madrone, myrtle, and bigleaf maple. The understory is huckleberry, salal, rhododendron, vine maple, willow swordfern, poison oak, and others. The major plant communities vary in age from early seral stages to old growth.

Natural climatic events have played a dominant role in shaping the stream channels in the watershed. Evidence from the 1955 and 1964 floods are contained on aerial photographs. Gravel bars were measured along the mainstem of Elk River on 1940-1986 aerial photos. The number of gravel bars on the whole watershed increased by 77% overall. In the upper segment which is wider and lower gradient, gravel bars increased more in size than in number. In the lower segment, which is narrower and steeper, a greater increase in the number of bars was observed.

Below the National Forest boundary, comparisons of the Elk River channel from 1940-1986 aerial photos show increased numbers and sizes of gravel bars. The unconfined channel flows through the valley floor show a change in pool geometry and frequency where new flood plains have been established over 46 years of flooding.

Elk River is recognized for its role in maintaining the viability of native salmonid stocks. The Elk River watershed produces anadromous steelhead trout, Coho salmon, Chinook salmon, and cutthroat trout. Resident rainbow and cutthroat trout populations are also present. Oregon Department of Fish and Wildlife operate a fish hatchery that was established in 1968 on the river just below Anvil Creek.

Adult Elk River Chinook salmon are characteristically three and four-year old fish that return primarily from November through January. This is a departure from many other coastal stocks, most of which return earlier in the fall. It is thought that the delay in adult spawning migration is an environmental adaptation to low-water conditions that persist on the southern Oregon coast during the fall months. The Elk River also appears to be the southern boundary between north-migrating and south-migrating coastal Chinook stocks (Maguire, 2001c).

Hubbard Creek

Hubbard Creek is located one mile southeast of Port Orford, Oregon. The watershed encompasses approximately 5340 acres or 8.34 square miles and is the primary water source for the City of Port Orford.

Basin Description:

- Boundaries: Elk River Watershed on the north and east.
- Garrison Lake Watershed on the north & west
- Rocky Point on the south
- Pacific Ocean on the southwest
- Elevation: Sea Level to 1200 feet

Description: Hubbard Creek is composed of three main branches, the north, middle, and south forks. The south fork is the longest having a reach of approximately 5 miles. The south fork joins the middle fork approximately 1 mile from the mouth, which then becomes the mainstem and the north fork joins the mainstem approximately one quarter mile from the mouth.

The north fork, at approximately 1 mile from the mainstem, has an impoundment, which serves as a reservoir for the city of Port Orford. This is the primary water source for this municipality (Maguire, 2001d).

Fish:

- North Fork – Cutthroat
- Middle Fork – Cutthroat
- South Fork – Cutthroat, Steelhead, Coho Salmon

Euchre Creek

Euchre Creek is only 14 miles long and drains from one of the smallest watersheds of any river in south coastal Oregon (23,831 acres/37 mi²). Elevations in the watershed range from sea level to approximately 3,080 ft. Major tributaries include Cedar and Boulder creeks. The upper portion of the basin is characterized by steeply sloped forested land with narrow valleys and tributary streams that have moderate to very steep gradients. The predominant land use in the middle and upper portions of the watershed is commercial timber production. The lower few miles of the river are relatively low gradient coastal floodplain. Rural residential development, grazing, and other agricultural uses are the dominant land uses in the lower basin.

Approximately 75% of the Euchre Creek watershed is located in the Southern Oregon Coastal Mountains ecoregion, which is characterized by complex geology; steep, high gradient streams; and seasonally abundant precipitation. Watersheds in this ecoregion tend to have high stream densities due to the potential for heavy precipitation during the winter months. Extensive erosion can result from fractured geology, high peak stream flow rates, and a high incidence of landslides in this ecoregion. The remaining 25% of the Euchre Creek drainage is located in the Coastal Uplands ecoregion, and is characterized by moderate to low gradient streams which are generally underlain

by sandstone, and flow through predominantly deep silt loam soils. Although peak precipitation rates are slightly lower than in the Southern Oregon Coastal Mountains, streams in the Coastal Uplands ecoregion are also susceptible to high erosion rates due to extreme fluctuations between summer and winter stream flows, easily eroded soils, and the prevalence of diked and channelized stream reaches. The risk ratings for increased erosion from roads within the Euchre Creek watershed ranged from moderately low to moderate.

Over three fourths of the land in the Euchre Creek basin is in private ownership, with almost 49% of the land owned and managed by private industrial timber companies, and 29% owned by small-parcel, private landowners. Public lands comprise almost 22% of the lands in the basin, with management responsibility for these lands split almost evenly between the Bureau of Land Management and the U.S. Forest Service. There are only 25 acres of State land in the Euchre Creek watershed.

Forestry is the most dominant land use in the Euchre Creek watershed, accounting for 94% of the total land area. The remaining 6% of the watershed is used for agriculture, animal range, and rural residential development. Rangelands are mostly managed for livestock grazing, with cows being the primary type of livestock grazed. To a lesser extent, sheep, llamas, goats, horses, and other small animals also use rangeland.

Riparian habitats in Euchre Creek are characterized by a scarcity of large conifers near the surveyed channels and a relatively low incidence of bank erosion. Shade is within or very close to the desirable range for all reaches. Aquatic habitat complexity is low, primarily because of the lack of large wood everywhere except Boulder Creek. Euchre Creek is not currently listed as water quality limited by the Oregon Department of Environmental Quality, but is being investigated for inclusion on the 303d list for temperature, flow modification, and sediment.

There are relatively few Out-of-Stream water rights for the Euchre Creek watershed. However, the majority of those rights are senior to the 1964 In-Stream water right of 10 cubic feet per second in the summer months. Out-of-Stream water rights currently exceed the flow in Euchre Creek from May to October of most years, with the majority of these water rights allocated for irrigation use. Water Storage Rights within the Euchre Creek watershed total 179 acre-feet, and are almost entirely allocated for industrial use. Most of these rights are not used but still active.

An estimated 90 acres of wetlands have been assessed in the Euchre Creek drainage. Most of the wetlands in the drainage are found in the lower gradient sections along the lower mainstem of Euchre Creek. Approximately 40 of these acres have been highly modified, 17 acres have been moderately modified, and the remaining 33 acres are not significantly altered. Wetlands within the Euchre Creek watershed are bordered primarily by agricultural land, but are bordered to a lesser extent by forested, rural residential, and other developed lands.

Chinook and Coho salmon, steelhead, and cutthroat trout are all native to the Euchre Creek drainage, although the historic abundance and distribution of these salmonids within the watershed are poorly understood (Maguire, 2001e).

Rogue River

The Rogue River basin extends from the west slope of Crater Lake on the east to the northernmost portion of California on the south to the Pacific Ocean on the west. The Basin covers 5,160 square miles and lies in the Klamath Mountain Province. The Rogue River is the third largest river in Oregon and is approximately 200 miles long. This Plan covers that portion of the basin that lies within Curry County.

Precipitation ranges from 80” per year at the coast to 130” at Signal Buttes. Before 1977 the annual discharge of the Rogue was 5.66 million acre-feet. After the completion of the Lost Creek Dam in 1977 and the Applegate Dam in 1980 the annual discharge fell to 3.97 million acre-feet. At the Agness gauge, the highest recorded flow was 290,000 cubic feet per second (cfs) in 1964 and the lowest was 608 cfs in 1968. The principal tributaries within Curry County are Shasta Costa Creek (river mile [RM] 29), Illinois River (RM 27), and Lobster Creek (RM 11).

The Rogue estuary extends from the ocean to Ferry Hole (RM 5) with several sloughs and multiple channels found in this reach. The substrate is primarily gravel with some areas of sand or mud bottom. There is almost no large woody debris present in the estuary because of the stream power. The river mouth and the Port of Gold Beach are protected by jetties constructed in 1960. Many wetlands and riparian terraces in the lower 15 miles of the river canyon have been converted to agriculture since 1850 when European settlers began arriving in the county.

The majority of the watershed consists of steep, forested slopes that are used for timber production. Grazing livestock on the meadows scattered through the forest was formerly common, but is now confined primarily to the coastal area. Ownership is primarily USFS in the interior with a fairly narrow strip of private lands along the coast and the river below Agness. Of that portion of the watershed between Agness and the mouth, 44,600 acres (54%) is USFS, and the remaining 38,000 acres are private (timber companies, rural residential, urban, and county lands).

A variety of anadromous and resident salmonids use the watershed including Chinook, winter and summer steelhead, Coho, cutthroat, and rainbow trout. The Rogue and Illinois mainstems are used as migration habitat while Lobster and Quosatana Creeks provide the majority of the spawning and rearing habitat. The smaller, steeper tributary streams provide cool water to help lower summer water temperatures in the mainstem. Other species utilizing the estuary and river include sturgeon (green and white), shad, and lamprey.

The human population of the Lower Rogue watershed is concentrated in the towns of Gold Beach (population 2000), Agness (population approximately 500), and in rural residential along the lower 10 miles of the Rogue and on Squaw Creek. The Rogue River continues to support a thriving recreational and commercial fishery as well as other aquatic activities (Weinhold, 1995).

Hunter Creek

The Hunter Creek watershed can be divided into five subwatersheds: Upper Hunter mainstem (North Fork confluence to headwaters), Middle Hunter mainstem (North Fork confluence down to Section 21), Lower Hunter mainstem (Section 21 to the mouth), North Fork Hunter, and Big South Fork Hunter. The whole watershed encompasses 44.4 square miles or 28,405 acres.

The watershed is divided between public (38%) and private lands (62%). The public lands are administered by the U.S. Forest Service, the Bureau of Land Management, and the state. Ownership of the private lands is dominated by approximately 81 % ownership by forest industry. Forestry uses account for 97% of the land in the watershed on both private and public lands. The remaining 3% of the watershed is divided between urban, agriculture/range and rural residential uses.

Precipitation ranges from 80” per year at the coast to 130” per year in the interior mountains. The majority of the watershed receives approximately 110” per year. Elevations range from sea level in the Lower Hunter mainstem sub-watershed to 3,558 feet on Sugarloaf Mountain in the Upper Hunter mainstem sub-watershed and 3,512 feet near Signal Buttes in the North Fork Hunter sub-watershed. The rain-on-snow zone encompasses 15% of the watershed - primarily in the Upper Hunter mainstem and North Fork Hunter sub-watersheds.

The majority of the watershed consists of steep, forested slopes that are used for timber production. Grazing livestock on the meadows scattered through the forest was formerly common, but is now confined primarily to the coastal valley area. Ownership is primarily USFS and BLM in the interior with the private timberlands owned by South Coast Lumber Company, Crook Estate, and Menasha Corporation.

A variety of anadromous and resident salmonids use the watershed including Chinook, winter steelhead, Coho, and cutthroat trout. The lower mainstem is used as migration habitat while the forks and tributaries provide the majority of the spawning and rearing habitat. The smaller, steeper tributary streams provide cool water to help lower summer water temperatures in the mainstem. Coho populations in Hunter Creek were smaller than Chinook due to the steep, confined high-energy streams. Cutthroat are thought to use most of the streams in the watershed, while steelhead use the lower section of each fork and the mainstem (Maguire, 2001f).

Pistol River

The Pistol River watershed drains approximately 67,275 acres or 105 square miles of land. Pistol River, situated entirely within Curry County, is an average size watershed on the southern Oregon coast. Flowing in a westerly direction Pistol River crosses Highway 101 and drains into the Pacific Ocean about ten miles south of the community of Gold Beach. Elevations in the watershed range from sea level to approximately 4,220 feet on Snow Camp Mountain. Major tributaries include the North Fork, East Fork, and South Fork. The upper portion of the watershed is characterized by steeply sloped forested areas with narrow valleys and tributary streams that have moderately steep to very steep gradient. Forestry is the dominant land use in the upper portion of the watershed

At one time there were six active lumber mills in the Pistol River area; a series of dairies in the lowlands; and a cheese factory. During the past forty year, some of these industries have left the area, but ranching, rural residential development, and other agricultural uses still thrive in the lower portion of the watershed. Over 55% of the watershed is in public ownership.

Southern Oregon Coastal Mountains make up 14% of the watershed with steep to very steep gradients, high rates of erosion, and high stream densities. Rainfall averages 79-140 inches per year.

The Coastal Siskiyou make up 82% of the watershed, with habitat very similar to Southern Oregon Coastal Mountains. Coastal Uplands cover less than 1 percent of the watershed and roughly follow the historic Sitka spruce distribution. High and low gradient habitats are present, with slow moving earth flows common on the hill slopes.

Approximately 177 acres of wetlands are found in this watershed. All the wetlands in the Pistol River watershed are in the Lower mainstem.

The Pistol River mainstem is listed on the 303(d) list for water temperature from the mouth to the headwaters. Temperatures range from the mid to high 60s in the mainstem Pistol River above East Fork, the East Fork, North Fork and Deep Creek. Temperatures range from high 60s to low 70s in the mainstem Pistol River above the South Fork, Crook Creek, and the South Fork Pistol. Temperatures range from mid to high 70s at the ODFW trap on the mainstem Pistol River.

Chinook distribution covers all of the mainstem Pistol River, up to 2/3 of Sunrise Creek, approximately half of the South Fork, and the lower portions of several mainstem Pistol River tributaries. No Chinook have been reported on the Upper Pistol and East Fork. Coho distribution is similar to Chinook with less use of the South Fork and no use of the Sunrise Creek. Steelhead

utilize nearly all of the mainstem, all of South Fork, one-third of the North Fork, one-third of the East Fork, and all three major mainstem tributaries (Maguire, 2001g).

Chetco River

The Chetco River is the second largest coastal watershed south of the Coquille drainage, and drains approximately 352 square miles, or 225,00 acres. It flows west out of the Siskiyou and empties south into the Pacific Ocean about 6 miles north of the California/Oregon state line between the cities of Brookings and Harbor, Oregon. The mainstem of the Chetco River is about 56 miles long, with the first 28 miles located within the Kalmiopsis Wilderness area. Elevations in the watershed range from 5,098 feet to sea level, and much of the watershed is characterized by steeply sloped forested valleys with moderately steep to very steep gradient. More than 80% of the watershed is public land, and the primary land manager is the United States Forest Service (USFS). Private land only occurs in the lowest 11 miles of the river, and the predominant land use is industrial and non-industrial forestry. Agriculture/grazing, rural residential development, and urban development together occupy less than 3% of the land use in the Chetco River drainage. The Chetco River estuary is short (< 2 miles in length) and is highly modified from its original condition by construction of a boat basin and jetties that extend into the Pacific Ocean.

Precipitation in the Chetco River watershed is lowest near the coast and increases to the north and east. Annual precipitation on the coast averages 75 inches, whereas the peaks in the Coast Range receive as much as 170 inches. Flows in the Chetco River are highly variable due to the seasonality of precipitation in southwestern Oregon and because there is little snowmelt to boost the river's flow in the spring and summer. Summer flows are often less than 100 cubic feet per second (cfs), whereas winter flows have exceeded 40,000 cfs in 5 of the last 30 years. Rain-on-snow events are common during the winter in the high steep mountains in the headwaters of the Chetco River, and can result in large day-to-day variability in flows. Due to minimal winter snowpack and the warm southwestern climate, the Chetco is consistently warmer than 65 degrees in the summer. DEQ modeling efforts confirm this.

Major fires historically occurred in the Klamath Mountains ecoregion on about 50 year intervals. Native Americans and ranchers both used fires to maintain grasslands and prairies. Prior to 2002, modern fire suppression policy had severely reduced the incidence of wildfire in this region leading to the 2002 Biscuit fire. Severe windstorms occur regularly in the Chetco basin, and can topple large patches of mature trees. High intensity rainfall events on steep slopes can result in many landslides and earth flows into the Chetco River.

Forestlands and their charred remains account for 97% of the Chetco watershed. Urban use accounts for almost 1% of the land use in the Chetco River basin, but is confined to the lower few miles of the river. The remaining 2% of the watershed is used by rural residential development, animal range, and a small amount of agriculture. Rangelands are managed for livestock grazing, whereas agricultural lands are primarily used for producing hay or gardening. The majority of the livestock on grazing lands are sheep and cattle. To a lesser extent, rangelands are also used by llamas, goats, horses, and other small animals.

The Chetco River has been placed on the 303d list because it exceeds the 64°F standard in three locations, one of which is just downstream from where the river emerges from the Kalmiopsis Wilderness area. Even though water from the Kalmiopsis wilderness exceeds the standard at that point, it cools as it moves toward the mouth. One of the factors for this cooling is the contribution of cooler water from downstream tributaries. The Chetco River is currently being investigated by the DEQ for additional 303d listings.

The out-of-stream water rights for the Chetco River total approximately 59 CFS. The majority (60%) of these water rights are allocated to municipal water use, and 73% of the rights are junior to

the in-stream water right (80 CFS) established by the Oregon Department of Fish and Wildlife for wildlife in 1964. Water storage rights total 370 acre feet in the Chetco watershed, with the majority of those rights allocated to municipal water storage. Out-of-stream water rights currently exceed flows in the Chetco River from July to October, and no additional water rights are available during those months. Water allocated for livestock and irrigation comprise less than 11% of the total out-of-stream water rights for the Chetco River.

An estimated 93 acres of wetlands have been assessed in the Chetco River watershed, with most of them found in the lower gradient sections along the lower mainstem, Jack Creek, and the North Fork of the Chetco River

Chinook and Coho salmon, steelhead, and cutthroat trout are all native to the Chetco River watershed. The historic abundance and distribution of these salmonids within the watershed is poorly understood. Historic numbers of Coho are thought to have been low in most south coast basins, including the Chetco River, due to relatively steep topography which results in high gradient, confined drainages which are not conducive to Coho spawning or rearing. Chinook salmon and steelhead are thought to occupy most of their historic habitat in the Chetco River watershed. Little is known about the historic distribution and abundance of cutthroat trout, but currently they are thought to fully utilize the available habitat. The Chetco River continues to support a healthy recreational and commercial fishery (Maguire, 2001h).

Winchuck River

The Winchuck River watershed is one of the smaller watersheds on the Southern Oregon coast. The Winchuck flows into the Pacific Ocean just north of the Oregon- California border. Its watershed is primarily within Curry County with some tributaries in California's Del Norte County (South Fork, Middle Winchuck mainstem, and upper Bear Creek).

The Winchuck River drains approximately 45,600 acres (71.4 square miles of land). Steep forested areas and narrow valleys characterize the upper portion of the basin. Approximately 70% of the Winchuck watershed is in public ownership (U.S. Forest Service and some State Parks).

The lower basin receives 50-70 inches of annual rainfall and the upper basin can receive 100-150 inches with winter snow accumulations. Rain-on-snow run-off occurs frequently and can have an influence on peak flow periods. Tidal movements affect the lower 1.5 miles of the Winchuck River. However, this river system has a very short estuarine system for juvenile salmonid rearing and exhibits sandbar closings at the mouth during late summer months.

The majority of the watershed is in timberland (95%) and is managed as a late successional reserve. The Winchuck system was heavily logged in the 1950's and 1960's. Present logging on private land includes Douglas fir thinning, alder management, and reversion back to fir stands.

Agricultural uses in the basin are livestock grazing (cattle) and some lily bulb production and are limited to the lower Winchuck River mainstem and South Fork. The agricultural uses involve less than 500 acres.

Rural residential areas account for less than 2% of the private lands and have been increasing in recent years. Water uses within the basin include residential, limited irrigation, livestock management, and instream uses (fish, wildlife, etc.). Residential landowners utilize groundwater wells, springs, and surface water holding tanks.

The Winchuck River system hosts an anadromous fish population that includes Chinook salmon, winter steelhead, coastal cutthroat trout, and Coho salmon. The winter steelhead runs and cutthroat trout stocks are considered healthy by American Fisheries Society standards. The Chinook salmon

population had been slowly decreasing since the 1970's, but in recent years has shown an increase (ODFW). Coho salmon populations have historically been limited and a small sustaining run exists in the South Fork watershed.

The Winchuck system is on the 303(d) list for temperature impairment and dissolved oxygen. This watershed has a characteristic that allows it to clear, after major storm events, faster than any river in Curry or Del Norte counties (including the Chetco and Smith Rivers).

Wetland areas in the lower Winchuck basin are estimated at 42 acres. Approximately 88% of these wetlands have a moderate to high degree of alteration with 61% located near residential development and 27% adjacent to agriculture (Maguire, 2001i).

Mission

Goal, Objective, Responsibility, and Intent

Goal of the Committee:

To set forth agricultural management opportunities that result in the continued protection of water quality in the watersheds of Curry County.

Objectives:

- To protect water quality by limiting, to the extent feasible, undesirable contributions from agricultural practices.
- To raise public awareness of agriculture's contribution in protecting water quality.
- To provide public involvement opportunities to share information about positive agricultural management practices.
- To have a plan that is developed locally, supported by the local people, implemented voluntarily, and which achieves regulatory water quality mandates for agricultural practices.

Responsibility of the Committee as stated in ODA Policy:

- OAR 603-090-0003 - Create an Agricultural Water Quality Management Area Plan that comprehensively outlines measures that will be taken to prevent and control pollution from agricultural activities...
- OAR 603-090-0024(b) - Recommend strategies necessary to achieve water quality goals and objectives...
- OAR 603-090-0030 - Describe a program to achieve water quality goals and standards necessary to protect beneficial uses related to water quality, as required by state and federal law. An Area Plan shall include, but not be limited to the following:
 - description of the geographic area to which the Area Plan applies
 - a listing of water quality issues of concern
 - a listing of current beneficial uses being adversely affected
 - a statement that the goal is to prevent and control water pollution from agricultural activities and to achieve water quality standards
 - a statement of water quality objectives of the Area Plan
 - a description of the pollution prevention and control measures deemed necessary to achieve the goal
 - a schedule for implementation adequate to meet dates described by law
 - guidelines for public participation
 - implementation and enforcement strategies

Water Quality Issues of Concern

Introduction

Agricultural presence in the Curry County is not new to the human population of this area. Cattle, dairy operations, and some row crops have been present in the Curry County basins for over 125 years. Today, actual agricultural acreage has dramatically declined and human populations and residential watershed impacts have increased. In no way does the following discussion single out agriculture as the sole contributor to water quality problems in Curry County. On the contrary, agricultural management can have a positive impact on water quality relative to many other potential land uses in these coastal basins.

We wish to make it clear that there are numerous natural, uncontrollable influences on the listed parameters within Curry County. Headwater streams coming out of the wilderness areas have periods when they are above the temperature standard for Oregon. Tidal influences, wildly variable stream flows, climatic variables, and geologic factors all influence surface water heating rates. It has been admitted by DEQ on several occasions that there are many streams in the state that will never meet the 64-degree standard. There is nothing agricultural operations, all in the lower part of the watersheds, can do to **meet** that standard. Instead, what we are asked to do by the state of Oregon is to reduce or eliminate our contribution to the rate of heating

Contributing Factors

As stated earlier, there are background water quality problems that are not due to human activities. Bacteria reside in streamside soils. Water temperature can be warmed by air temperatures and stream channel substrate. Sediment and bank erosion are part of the natural hydrologic and geologic system. Nutrients, such as phosphorus, can be dissolved from parent rock material. Background sources of pollutants can be very hard and costly to identify and distinguish from management related sources.

Population increases and their resulting environmental impacts have changed the face of several stream systems within Curry County. Changes in fire frequency, the severity of peak and low stream flows, waste inputs, flood plain encroachment, degraded riparian areas, and airborne pollutants are all consequences of human population expansion into aquatic and terrestrial habitat. These are consequences that can be buffered but never eliminated. We are now an integral part of the landscape and environment.

The following narrative, tables, and lists focus on the mandate of the SB1010 legislation. Agriculture activities are only a small part of the land use in these basins. The conditions identified by the farmers and ranchers of the Local Advisory Committee will meet the stewardship and conservation needs on private agriculture lands to help alleviate the cumulative effects of our human impacts in all Curry County basins.

303(d) list -Addressing 303(d) listed Parameters

As of 2004, the most prevalent water quality issue in Curry County is the exceedence of the 64°F water temperature criterion. Table 1 below illustrates the percentage of the Curry County Basins stream segments listed on the 303(d) list that exceed the federal Clean Water Act standards for temperature and other parameters for which valid data sets are available. **These are NOT listed because of agricultural activity in the basin. In many cases stream segments are listed as temperature limited without any human induced factors present at all.**

Table 1

Parameter (from 2002 303(d) list)	Stream Segments Exceeding Standard
Temperature [64° F average daily maximum]	26/27 = 96%
Dissolved Oxygen [less than 90% saturation or 11 mg/l or 8.0 mg/l intergravel]	3/27 = 11%
pH [6.5-8.5 range]	1/27 = 4%
Aquatic Weeds or Algae [...having a deleterious effect on stream bottoms, fish or other aquatic life or ...injurious to health recreation or industry...]	1/27 = 4%

The Coastal Zone Act

On November 5, 1990, Congress enacted the Coastal Zone Act Reauthorization Amendments (CZARA). This law mandated that all states and territories with approved coastal zone management programs develop and implement coastal nonpoint source pollution control programs. In response to these CZARA Amendments, Oregon identified coastal SB 1010 plans as the state's strategy to address certain agricultural measures. The exact text of the approved Management Measures is included in Appendix B. The CZARA management measures provide guidance when developing a plan for streams in the coastal zone of Oregon.

Prevention and Control Measures

The focus of the Agricultural Water Quality Management Program is on voluntary and cooperative efforts by landowners and others to protect water quality. However, Senate Bill 1010 also provides for a regulatory backstop to ensure prevention and control of water pollution from agricultural sources in cases where landowners or operators refuse to correct problem conditions. Agricultural Water Quality Management Area Rules serve as this backstop while allowing landowners flexibility in how they protect water quality. Area Rules are goal-oriented and describe characteristics that should be achieved on agricultural lands, rather than practices that must be implemented.

ODA appointed the Local Advisory Committee to assist in developing Area Rules (Prevention and Control Measures) to protect water quality and prevent and control water pollution from agriculture. On an 8 to 4 vote, the LAC submitted a riparian rule to ODA. The rule submitted was not consistent with legislative direction, so it was left to ODA to develop the rules that apply to the Curry Agricultural Water Quality Management Area. To do this, ODA looked to work developed by other LACs in similar planning areas for a model rule to address riparian conditions. The rules listed below are modified to reflect the unique circumstances found in Curry County.

OAR 603-095-3540

(1) All landowners or operators conducting activities on lands in agricultural use must comply with the following criteria. A landowner is responsible for only those conditions resulting from activities controlled by the landowner. A landowner is not responsible for conditions resulting from actions by another landowner on other lands. A landowner is not responsible for conditions resulting from unusual weather events or other exceptional circumstances that could not have been reasonably anticipated. A landowner is not responsible for natural increases in nutrient or temperature loading.

Definitions

Oregon Administrative Rule (OAR) 603-095-0010(4) defines agricultural use as “the use of land for the raising or production of livestock or livestock products, poultry or poultry products, milk or milk products, fur-bearing animals, or for the growing of crops such as, but not limited to, grains, small grains, fruit, vegetables, forage grains, nursery stock, Christmas trees, or any other agricultural or horticultural use or animal husbandry or any combination thereof. Wetlands, pasture, and woodlands accompanying land in agricultural use are also defined as in agricultural use.”

The following rules apply to any agricultural use exceeding 10,000 square feet in area, including, but not limited to tilling, clearing, grading, excavating, grazing, and feedlot usage; and excluding minor land disturbing activities such as home gardens and individual landscaping and maintenance (OAR 603-095-0010(43)).

Statutes Addressing Water Pollution

In 1995, the Oregon legislature recognized potential confusing authorities that belonged to both ODA and DEQ regarding the enforcement of water quality statutes. To clarify authorities granted to ODA in Senate Bill 1010, the state legislature passed Senate Bill 502, which was codified into ORS 561.191. This statute states that ODA shall develop and implement any program or rules that directly regulate farming practices that are for the purpose of protecting water quality. A 1996 opinion from the Oregon Attorney General's office states that **ODA has the statutory responsibility to regulate agriculturally related water pollution**. That same opinion also recognized the need to define that authority by developing water quality plans and rules that specifically address agricultural practices and land conditions and achieve the standards adopted by the Environmental Quality Commission

To implement Senate Bill 502, the Department is incorporating ORS 468B.025 and ORS 468B.050 into all of the basin Agricultural Water Quality Management Administrative Rules in the state. ORS 468B.025 and ORS 468B.050 are being incorporated by including the following language in individual basin administrative rules:

OAR 603-95-3540

(2) Effective on rule adoption, no person subject to these rules shall violate any provision of ORS 468B.025 or ORS 468B.050.

Upon adoption of this Rule, ODA assumes responsibility for implementing ORS 468B.025 and 468B.050.

This rule references current State Law (ORS 468B.025 and ORS468B.050). ORS 468B.025 states that no person shall:

- (1)(a) Cause pollution of any waters of the state or place or cause to be placed any wastes in a location where such wastes are likely to escape or be carried into the waters of the state by any means.
 - (b) Discharge any wastes into the waters of the state if the discharge reduces the quality of such waters below the water quality standards established by rule for such waters by the Environmental Quality Commission.
- (2) Violate the conditions of any waste discharge permit issued under ORS 468B or ORS 568.

ORS 468B.050 refers to situations when permits are required, such as for certain confined animal feeding operations.

Compliance with this rule ensures that concentrated nutrients, pathogens associated with high animal density areas, high sediment concentrations in run-off, toxics, or other potential pollutants are not readily transported to waters of the state.

Livestock wastes can include manure from pastures draining to or bisected by irrigation ditches and any other situations not already covered by Oregon's Confined Animal Feeding Operation laws. Indicators of potential noncompliance include: 1) runoff flowing through areas of livestock usage and entering waters of the state, 2) livestock waste located in drainage ditches or areas of flooding, or 3) E. coli counts that exceed State water quality standards. Livestock facilities located near streams should employ an adequate runoff control and waste management system.

Wastes can also include *excess sediment discharges*. Indicators of potential noncompliance with Rule (3) include: 1) visible active erosion scars, 2) sediment-laden runoff, or 3) obvious deposits of sediment on the stream or canal bottom that can be traced to a specific source.

Definitions:

Wastes include manure, commercial fertilizers, soil amendments, composts, vegetative materials, or *any other substances* that will or may cause water pollution (ODA's OAR 603-095-0010(53)). Therefore, 'wastes' also include sediment.

Waste discharge means the discharge of waste, either directly or indirectly, into waters of the state (ODA's OAR 603-095-0010(54)).

Water pollution means such alteration of the physical, chemical or biological properties of any waters of the state, including change in temperature, taste, color, turbidity, silt or odor of the waters, or such discharge of any liquid, gaseous, solid, radioactive or other substance into any waters of the

state, which will or tends to, either by itself or in connection with any other substance, create a public nuisance or which will or tends to render such waters harmful, detrimental or injurious to public health, safety or welfare, or to domestic, commercial, industrial, agricultural, recreational or other legitimate beneficial uses or to livestock, wildlife, fish or other aquatic life or the habitat thereof (State Statute for Water Quality: ORS 468B.005(7)).

Waters of the State include lakes, bays, ponds, impounding reservoirs, springs, wells, rivers, streams, creeks, marshes, inlets, canals, and all other bodies of surface or underground waters, natural or artificial, public or private (except those private waters which do not connect to natural surface or underground waters) within Oregon (from State Statute for Water Quality: ORS 468B.005(8)).

Riparian Areas

OAR 603-95-3540

(3) Effective June 3, 2007, agricultural management activities in the riparian area of perennial streams will be conducted in a manner that allows for the establishment, growth, and maintenance of riparian vegetation consistent with vegetative site capability so as to provide streambank stability and shade.

(a) Exemptions from OAR 603-095-3540(3) are:

(A) Stream crossings, access for irrigation equipment and other accepted water dependent agricultural uses when conducted in a manner that minimizes impacts on streambank stability

(B) Streams that do not support native trout and are inaccessible to anadromous fish because of barriers at their junction with the Pacific Ocean.

(C) This rule is not intended to prohibit riparian grazing where it can be done while meeting the above vegetative conditions.

This rule was developed to clearly show that landowners are not required to have mature riparian vegetation immediately but only that they must allow for the establishment, growth and maintenance. It does not require that adequate riparian vegetation be present by 2007. For guidance on management activities that promote the growth and establishment of riparian vegetation, please contact the Curry SWCD. For more information on the effects of riparian vegetation on water quality, please consult the References section.

This rule specifies that “agricultural activities” must allow for riparian vegetation to develop to make it clear that landowners are not responsible for the impacts of browsing activities of wildlife such as elk, geese, and beaver.

Definitions:

A riparian area is an edge or bank of a river, tributary, or other body of water.

“Site capability” means the ability of a site to provide for the development of potential structural and functional properties. Structural properties include, among other things, vegetation and soil characteristics. Functional properties include processes such as energy and nutrient flow. Capabilities to produce and sustain these properties are site-specific (Barrington et al, 2001). For additional references on site capability and related concepts, please consult the following in the references section: Gregory et al, 1991; Hunsaker and Levine, 1995; Leonard et al, 1992; Montgomery, 1999; Palik et al, 2000; Prichard, 1998; Winter, 2001; Winward, 2000.

ODA is responsible for determining site capability. This determination is based on soils, topography, climate, and other site characteristics that are described in U.S. Department of Agriculture soil surveys and through on-site visits.

Pesticides

Pesticide control is presently regulated by ODA under ORS 634 and OARs 603-057. Waterbodies in this basin have not been identified on the 3030(d) list for pesticide contamination. Thus, potential pesticide issues are addressed under existing statute and no additional measures are needed.

Strategies for Implementation

The Curry Soil and Water Conservation District can provide direct assistance to landowners in developing Area Plans for individual landowners. The Curry SWCD, as the designated local management agency, will provide specific landowner education and support for this Area Plan. Currently, the following activities are proposed by the SWCD:

- Regular SWCD newsletter; containing articles on positive management practices, outstanding District cooperators and their management practices, and articles about the status of the Curry County AgWQM Area Plan and Rules;
- Workshops, presentations, and seminars that will relate to the unacceptable conditions and positive management practices that are found in the Curry County AgWQM Area Plan;
- Press releases, and meeting announcements concerning Curry Area activities;
- Coordination between other agencies and associations such as DEQ, OSU Extension, Watershed Associations, etc.;
- Serve as a general clearinghouse of information for the public about SB 1010 and also an archive for the Curry LAC;
- Provide incentive for the development of Voluntary Farm Plans in the Curry Area;
- Employment of a farm planner(s) and other staff to facilitate the implementation of SB 1010 in the Curry Area.

This Area Plan in draft form was presented and made available to the public for public comment. Testimony presented at public hearings and collected during public comment periods were reviewed by the ODA and the Local Advisory Committee, and recommended modifications to the AgWQM Area Plan were presented to the Board of Agriculture and the director of the ODA for their review and comment. The final OARs resulting from this process were adopted through the Administrative Rules process by the director of the ODA.

This Area Plan and the associated Area Rules are subject to a two-year review process. Two years after adoption, ODA, in cooperation with the Curry SWCD and the Curry Local Advisory Committee, will assess the progress of Area Plan implementation toward achievement of Area Plan goals and objectives. Any new water quality information and programs, including Total Maximum Daily Loads, affecting agriculture will be reviewed and considered during this review process. As needed, recommendations to the State Board of Agriculture and director will be made regarding modifications to the Area Plan that may be necessary to achieve water quality goals and objectives. Any future amendments to the administrative rules will be subject to public participation process as defined in Oregon law.

Monitoring and Evaluation

Monitoring is an important activity as part of the implementation phase of Agricultural Water Quality Management (AgWQM) Area Plans. When effectively used, monitoring and data analysis can provide valuable information to:

- Establish baseline information;
- Evaluate trends in water quality improvement;
- Help understand whether water quality improvement activities are achieving their intended goals; and
- Assist with adjustments in implementation activities and priorities to gain maximum effects on improving water quality and watershed conditions.

Landowners interested in monitoring can find help locally through OSU, the Curry SWCD, the Curry Watershed Associations, DEQ, and other public and private sources.

Monitoring and assessment are also important information gathering activities during site-specific determinations of compliance as part of an investigation. For the purposes of AgWQM Area Plans, there are four important types of monitoring or assessment that may be conducted.

Baseline Condition Monitoring

Baseline condition monitoring provides a starting point for assessing water quality trends and for future evaluation of the effectiveness of water quality improvement efforts. Baseline condition monitoring typically includes identification and analysis of data previously and currently collected in the area according to accepted protocols. The Oregon Plan Water Quality Monitoring Technical Guide Book is the recommended guide for baseline condition monitoring.

Water Quality Trend Monitoring

Water quality trend monitoring can help to track how water quality (typically on a watershed or subwatershed scale) is changing over time, including after implementation of an AgWQM Area Plan. It is recommended that trend monitoring follow recommendations in the Oregon Plan Water Quality Monitoring Technical Guide. This Water Quality Monitoring Technical Guide Book describes accepted procedures and protocols for most activities that would be used to conduct baseline condition and trend monitoring on a watershed scale, including development of quality assurance/quality control plans to assure quality of data. Protocols described in the Water Quality Monitoring Technical Guide Book meet DEQ standards for data collection.

Effectiveness Monitoring

Effectiveness monitoring can be used to:

- a) Evaluate the effectiveness of specific management practices in reducing losses or loadings of components such as sediment or nutrients. The Natural Resources Conservation Service has a good amount of information about the effectiveness of various practices in protecting surface and groundwater quality.
- b) Evaluate the net effect of the implementation of an AgWQM Area Plan and watershed improvement activities on water quality trends.

Site-Specific Rule Compliance Monitoring and Assessment

Conducted as a part of a compliance investigation, this type of monitoring is specific to individual sites. It is performed to assess compliance with conditions in a rule, and to assess the contribution of land management activities or land conditions to rule or standards violations attributable to the landowner's activities. Site-specific information and data is collected to characterize and quantify the physical setting and land management conditions that relate to a potential rule or standards violation. Photographic documentation of the suspected problem is typically also included in the assessment. Water samples may be taken for analyses.

Current and future monitoring and assessment efforts

Monitoring of various water quality parameters is presently being done by several entities, including the Watershed Associations, the Department of Environmental Quality, the Oregon Department of Fish and Wildlife, and individual landowners. The data from these monitoring efforts can be analyzed and used for baseline condition monitoring, water quality trend monitoring, and effectiveness monitoring.

When implementation of this plan begins, ODA, in cooperation with other entities conducting monitoring activities, will pursue a monitoring strategy that will include an evaluation of baseline conditions. The department will not duplicate existing monitoring efforts being conducted by other

agencies or entities. The monitoring efforts that are pursued will be adapted to the changing conditions and issues that develop as the plan is being implemented.

Education and Outreach

The Oregon Department of Agriculture will coordinate the development of SB 1010 education projects within the Curry County with the Curry Soil and Water Conservation District. They will work hand in hand with US Department of Agriculture's Natural Resource Conservation Service, the OSU Extension Service, and the Curry County Watershed Associations to carry out an effective water quality education program.

To define, implement, and measure the success of the Curry County education effort, the following quantifiable tasks can be pursued:

1. Conduct education programs to promote public awareness of water quality issues.
 - Hold workshops on water quality issues and the conservation practices that will help improve water quality.
 - Develop demonstration projects to highlight successful conservation practices and systems.
 - Organize tours of demonstration projects for agricultural managers and producers.
 - Produce and distribute brochures about water quality issues.
 - Prepare standard presentations for agricultural producer groups.
 - Develop detailed, one-page Curry County fact sheets for erosion control, nutrient and waste management, livestock and grazing management, and riparian and streambank management.
 - Conduct one-on-one and small group visits with landowners to discuss the Curry County AgWQM Area Plan and adaptive management solutions.
2. Conduct a media program to inform Curry County agricultural operators, rural landowners, and the public of conservation issues and events.
 - Submit news articles and public service announcements to area newspapers, radio stations, and newsletters.
 - Invite media to conservation tours and workshops.
 - Include updates on the status of the Curry County AgWQM Area Plan and water quality data in Curry County SWCD, OSU Extension, and watershed council newsletters.
3. Involve the agricultural community in conservation education.
 - Create and maintain a list of experienced agricultural operators willing to share management solutions with other interested people by speaking, leading tours, and providing tour sites.
4. Build partnerships with commodity groups to promote conservation.
 - Co-sponsor workshops and tours among the commodity groups., watershed councils, and the Curry County SWCD
 - Share education materials with commodity groups and their representatives.
 - Develop educational materials in conjunction with commodity groups and watershed councils.
 - Partner with other agricultural and natural resource agencies, watershed councils, and commodity groups to access and acquire the material and financial resources to implement the Curry County AgWQM Area Plan and its educational component.
 - Meet with other agencies and organizations, and develop a strategy to obtain funding from traditional and nontraditional sources.

We believe that the vast majority of landowners want to do the things that will benefit the land and their production. A great deal of effort and resources should be used to inform landowners of the management strategies that will improve both their land and the quality of their water.

Administrative Roles and Responsibilities

Total Maximum Daily Loads

The Oregon Department of Environmental Quality, in accordance with the Federal Clean Water Act, is required to establish Total Maximum Daily Loads (TMDLs) for waterbodies on the 303(d) list. TMDLs will identify the maximum amount (load) of each pollutant that a listed waterbody can receive and still meet state water quality standards. Once a Total Maximum Daily Load is established for a particular pollutant, each source of pollution in the area will be assigned a portion of that load, and each source must develop or modify pollution control plans and programs designed to achieve their load.

TMDLs are scheduled to be completed for Curry County in 2005. The Area Plan and Rules seeks to satisfy agriculture's load in the TMDLs for these waterbodies. Once TMDLs are completed for the Management Area, ODA and DEQ will compare the TMDLs and Area Plan and Rules to determine if any adjustments need to be made to the Area Plan and Rules to achieve agriculture's load allocations.

Designated Management Agency/Local Management Agency

The Oregon Department of Agriculture is the "Designated Management Agency" for nonpoint source pollution control activities on agricultural and rural lands in the Curry County basins. In turn, through Memoranda of Agreement, ODA has designated the Curry Soil and Water Conservation District as its Local Management Agency to assist with the development and implementation of the water quality management area plan and projects in the South Coast Basin. Implementation priorities will be established on a periodic basis through annual work plans developed jointly by the Curry SWCD and ODA, depending on available funding.

The Director of the Oregon Department of Agriculture appointed a Curry County Local Advisory Committee representing local agricultural producers, local landowners, local environmental interests, and the Curry SWCD for the purpose of assisting with the development of this Area Plan and the associated draft Oregon Administrative Rules to implement core elements of the plan. The draft Area Plan and Rules have been presented to the Oregon Board of Agriculture for their review and consultation.

The Local Management Agency and the Local Advisory Committee will participate in biennial review of plan implementation progress. Any future amendments to the administrative rules will be subject to the public participation process outlined in Oregon law.

Monitoring and Evaluation of the Plan's Effectiveness

The progress and success of implementation efforts will be assessed through determination of changes in land management systems and the measurement of water quality improvement over time. The number of private and public groups doing water quality trend monitoring will insure the LAC's awareness of water quality trends throughout the basin. ODA plans to conduct land condition assessments and outreach evaluations but will leave water quality monitoring to those who are funded for that task.

Two years after Area Plan and Rule adoption, the ODA, with the cooperation and assistance of the Curry SWCD, the LAC, and the DEQ, will assess the progress of plan implementation toward achieving plan's goals and objectives. These assessments may include:

1. An accounting of the number of operations with approved farm plans that address water quality concerns, the number of requests for farm plans, and the acreage they cover.

2. An estimate of the amount of pollution prevented based on number of farm plans implemented and their acreage.
3. Identification of additional agricultural sources of sediment, nutrients, and other contributors to 303(d) listed streams not addressed in the original plan.
4. An evaluation of the effectiveness of outreach and education programs designed to provide public awareness and understanding of water quality issues.
5. A review of projects, demonstrations, and tours used to showcase successful management practices and systems.
6. An evaluation of the effectiveness of the sources for technical and financial assistance that are available to the agricultural community.
7. Review of load allocations as found in Curry Subbasin's TMDL and the effectiveness of this plan in meeting load allocations.

Complaints and Inspections

It is the intent of this plan that the Oregon Department of Agriculture consider severe weather conditions when evaluating compliance with unacceptable conditions adopted in the rules. The department recognizes that every farm and situation is different and will take into account each individual situation when evaluating compliance with unacceptable conditions listed in the rules.

Compliance Procedures

Written complaints against operators or landowners alleged to be out of compliance with the rules associated with this Area Plan will be investigated by the Oregon Department of Agriculture. The complaint must identify the site, the specific information related to the undesirable condition, and be filed with the department. Copies of the complaint are available to the party alleged to be in violation. Based on the investigation, the department will determine if a condition violation exists and describe appropriate action to remedy the condition that violates the associated rules.

If an approved farm plan which addresses water quality concerns exists for a site which has been determined to have a conditions violation, and if the plan is being implemented on schedule, the operator and/or landowner will be given an opportunity to refine or modify the plan, or to develop an updated implementation schedule to remedy the condition within a specified time frame. The revised farm plan and/or implementation schedule must be approved by the LMA to verify its compliance with NRCS/FOTG standards and Area Rules, whichever is applicable.

When a Notice of Noncompliance is issued for an AgWQM Area Rule conditions violation, the landowner/operator will be directed by the Oregon Department of Agriculture to remedy the condition under provisions in OAR 603-090-0060 through 603-090-0120. **It is the policy of the department to direct the landowner to available assistance through state and federal partners so as to achieve a resolution rather than an enforcement action.** If an approved farm plan does not exist for the site, the operator and/or landowner can be given assistance to create and implement a farm plan. A farm plan, however, is not necessary to be in compliance with the rules. Authority for any enforcement action rests with the Oregon Department of Agriculture.

Review Period

The Oregon Department of Agriculture, Curry SWCD, the Local Advisory Committee, and the Oregon Board of Agriculture will consider making appropriate modifications to the basin plan and/or the associated Oregon Administrative Rules on a biennial basis.

The Curry Agricultural Water Quality Management Local Advisory Committee will be asked to reconvene two years after adoption of this plan. The purpose of reconvening is to address the implementation, language, or newly listed 303(d) parameters of concern to agriculture in the basin.

For the biennial review, ODA, and the Curry SWCD, propose to conduct a random survey of a statistically significant number of anonymous agricultural holdings in the Curry/South Coast Basin. This survey will establish a baseline of compliance with OAR 603-95-3840 upon which the LAC can reconvene to set benchmark compliance targets. The benchmarks will not have any regulatory impact. They are only to be used as guidelines for prioritizing educational activities and incentives.

References

- Barrington, M., D. Wolf and K. Diebel. 2001. Analyzing Riparian Site Capability and Management Options. *Journal of the American Water Resources Association* 37(6):1665-1678.
- Buzzard, C. and C. Bowsby. 1970. Soil Survey of the Curry Area, Oregon. U.S. Department of Agriculture, Washington, D.C.
- Gregory, S. V., F. J. Swanson, W. A. McKee, and K. W. Cummins. 1991. An Ecosystem Perspective of Riparian Zones. *Bioscience* 41(8):540-551.
- Hunsaker, C.T. and D.A. Levine. 1995. Hierarchical Approaches to the Study of Water Quality in Rivers. *Bioscience* 45(3): 193-203.
- Leonard, S. G., G. J. Staidl, K. A. Gebhardt, and D. E. Prichard. 1992. Viewpoint: Range Site/Ecological Site Information Requirements for Classification of Riverine Riparian Ecosystems. *Journal of Range Management* 45:431-435.
- Maguire, C. 2001a. Floras Creek Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001b. Sixes River Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001c. Elk River Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001d. Port Orford Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001e. Euchre Creek Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001f. Hunter Creek Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001g. Pistol River Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001h. Chetco River Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Ibid. 2001i. Winchuck River Watershed Assessment. South Coast Watershed Council, Gold Beach, Oregon.
- Massengill, C. 2003. Coastal Oregon Riparian Silviculture Guide. Coos Watershed Association, Charleston, Oregon.
- Montgomery, D. R. 1999. Process Domains and the River Continuum. *Journal of the American Water Resources Association* 35(2):397-411.
- Natural Resources Conservation Service. 1997a. Conservation Practice Standard for Tree and Shrub Establishment. Natural Resources Conservation Service, Portland, Oregon.

- Ibid. 1997b. Conservation Practice Standard for Riparian Forest Buffer. Natural Resources Conservation Service, Portland, Oregon.
- Ibid. 1997c. Conservation Practice Standard for Fence. Natural Resources Conservation Service, Portland, Oregon.
- Ibid. 1997d. Conservation Practice Standard for Nutrient Management. Natural Resources Conservation Service, Portland, Oregon.
- Ibid. 1997e. Conservation Practice Standard for Pest Management. Natural Resources Conservation Service, Portland, Oregon.
- Ibid. 1997f. Conservation Practice Standard for Livestock Watering Facility. Natural Resources Conservation Service, Portland, Oregon.
- Ibid. 1997g. Conservation Practice Standard for Use Exclusion. Natural Resources Conservation Service, Portland, Oregon.
- Ibid. 1997h. Conservation Practice Standard for Irrigation Water Management. Natural Resources Conservation Service, Portland, Oregon.
- Oregon Department of Agriculture. 2000. Relationship Between Agricultural Water Quality Management Area Plan Conditions and Water Quality Standards. Oregon Department of Agriculture, Salem, Oregon.
- Ibid. 2000. Relationship Between Agricultural Water Quality Management Area Plan Conditions and Water Quality Standards Part 2: Specific Research on Water Quality Impacts from Agriculture in Oregon. Oregon Department of Agriculture, Salem, Oregon.
- Palik, B.J., P.C. Goebel, L.K. Kirkman, and L. West. 2000. Using Landscape Hierarchies to Guide Restoration of Disturbed Ecosystems. *Ecological Applications* 10(1):189-202.
- Prichard, D. 1998. Riparian Area Management: A User Guide to Assessing Proper Functioning Condition and Supporting Science for Lotic Areas. USDI/BLM Technical Reference 1737-15. Bureau of Land Management, Denver, Colorado.
- Weinhold, M. 1995. Lower Rogue River Basin Watershed Condition Assessment. Lower Rogue Watershed Council, Gold Beach, Oregon.
- Winter, T.C. 2001. The Concept of Hydrological Landscapes. *Journal of the American Water Resources Association* 37(2):335-349.
- Winward, A. H. 2000. Monitoring the Vegetation Resources in Riparian Areas. USDA/US Forest Service General Technical Report RMRS-GTR-47. Rocky Mountain Research Station, Ogden, Utah.

Appendix A. Available Technical and Financial Assistance

Many agricultural landowners are unable to make a living directly from their land-based enterprise. Financial incentives are essential to encourage basinwide adoption of sound and sustainable management practices.

- CREP - Conservation Reserve Enhancement Program (541-396-4323) made available \$250 million dollars to pay landowners to set aside areas immediately adjacent to anadromous fish-bearing streams. It is intended to protect water quality and enhance spawning, rearing, and habitat quality.
- OWEB - Oregon Watershed Enhancement Program (541-471-2886) provides funding for watershed enhancement projects under the general categories of education/public awareness, monitoring, management, and assessment/action planning.
- EQIP - Environmental Quality Incentives Program (541-396-4323) pays landowners a majority cost-share for on-farm projects that protect natural resources and improve wildlife (including fish) habitat. EQIP information can also be obtained from the Farm Service Agency.
- EPA 319 - Environmental Protection Agency administers the 1972 Clean Water Act section 319 grants through ODEQ (541-269-2721x27) to help meet their water quality mandates. The projects EPA likes to fund are those with directly measurable benefits for water quality and endangered species. Check out EPA's Ag Info Center: <http://es.epa.gov/oeca/ag/index.html>
- NRCS - Natural Resources Conservation Service (541-396-4323) can provide technical assistance and administers a number of cost-share programs for on-farm projects that improve farm production while protecting natural resources and improving wildlife (including fish) habitat.
- OWT - Oregon Water Trust (503-226-3480 in Portland) offers lease and buy-out options for abandoned or unused water rights. This market-based approach to increasing stream flow may also be used to fund irrigation system changes in watersheds identified as priorities for OWT.
- OSU Cooperative Extension (1-800-356-3986 in Curry County) offers a wide variety of levels of technical assistance and planning help. OSU has been instrumental in the Oregon Cattlemen's extremely successful Watershed Ecosystem Education Program. Since its inception, it has grown into several distinct natural resource related workshops that are offered to ranchers and farmers free of charge. The Watershed Ecosystem Education Program workshops help ranchers and farmers understand their watersheds and stream function better through assessments and monitoring. OSU has also been providing Proper Functioning Condition (PFC) Workshops and assessments with landowners. PFC assessment should be a major component of a conservation plan.

Watershed Councils and Soil and Water Conservation Districts are a primary resource for finding technical and financial assistance. Contact the Curry SWCD at 541-274-2755.

Curry County hosts several watershed councils unique to their basins of concern. A call to the Curry SWCD can give you contact information for those unique watershed councils.

Appendix B. Coastal Zone Management Act Measures

In 1990, the Federal Coastal Zone Reauthorization Amendments were enacted. This law mandated that all states and territories with approved coastal zone management programs develop and implement coastal nonpoint pollution control programs. Listed below are the Coastal Zone Management measures that were developed for use in Oregon for coastal basins such as Curry County.

The following section contains the approved management measures for coastal nonpoint pollution in Oregon as developed for the Coastal Zone Reauthorization Amendments. For more detailed information on how to implement any of the recommendations below, consult one of the technical groups in Appendix A or publications included in the References section.

Sedimentation

- Apply the erosion component of a Resource Management System as defined in the Field Office Technical Guide of the U.S. Department of Agriculture, Natural Resources Conservation Service to minimize the delivery of sediment to surface waters.
- Design and install a combination of management and physical practices to settle the settleable solids and associated pollutants in runoff delivered from the contributing area for storms of up to and including a 10 year, 24 hour frequency.

Nutrients

- Develop, implement, and periodically update a nutrient management plan to: (1) apply nutrients at rates necessary to achieve realistic crop yields, (2) improve the timing of nutrient application, and (3) use agronomic crop production technology to increase nutrient use efficiency. When the source of the nutrients is other than commercial fertilizer, determine the nutrient value and the rate of availability of the nutrients. Determine and credit the nitrogen contribution of any legume crop. Soil and plant tissue testing should be used routinely.

Pesticides

- Evaluate the pest problems, previous pest management practices, and cropping history.
- Evaluate the soil and physical characteristics of the site, including mixing, loading, and storage areas for potential of leaching or runoff of pesticides. If leaching or runoff is found, steps should be taken to prevent further contamination
- User integrated pest management (IPM) strategies that:
 - Apply pesticides only when an economic benefit to the producer will be achieved (i.e. application based on economic thresholds).
 - Apply pesticides efficiently and at times when runoff losses are unlikely.
 - When pesticide applications are necessary and a choice of registered materials exists, consider the persistence, toxicity, runoff potential, and leaching potential of products being used.
 - Periodically calibrate pesticide-spraying equipment.
 - Use anti-backflow devices on hoses used for filling tank mixtures.

Riparian Areas

- Exclude livestock from riparian areas that are susceptible to overgrazing and when there is no other practical way to protect the riparian area when grazing uplands.
- Provide stream crossings and hardened access areas for watering.
- Provide alternative drinking water locations.
- Locate salt and shade away from sensitive riparian locations.
- Include riparian areas in separate pastures with separate management objectives and strategies.
- Fence, or where appropriate, herd livestock out of areas for as long as necessary to allow vegetation and streambanks to recover.
- Control the timing of grazing to: (1) keep livestock off streambanks where they are most vulnerable to damage, and (2) coincide with the physiological needs of target plant species.

Irrigation

- Operate the irrigation system so that the timing and amount of water match crop water needs. This will require, at a minimum: (a) the accurate measure of soil water depletion and the volume of irrigation applied, and (b) uniform application of water.
- When chemigation is used, include backflow preventers for wells, minimize the harmful amounts of chemigated waters from the field, and control deep percolation.
- In cases where chemigation is performed with furrow irrigation systems, a tailwater management system may be needed.
- In some locations, irrigation return flows are subject to other water rights or are required to maintain stream flow(s). In these special cases, on-site use could be precluded and would not be considered part of the management measures for such locations.
- In some locations, leaching is necessary to control salt in the soil profile. Leaching for salt control should be limited to the leaching requirement for the root zone.
- Where leakage from delivery systems or return flows support wetlands or wildlife refuges, it may be preferable to modify the system to achieve a high level of efficiency and then divert the “saved water” to the wetland or wildlife refuge. This will improve the quality of water delivered to wetlands or wildlife refuges by preventing the introduction of pollutants from irrigated lands to such diverted water.
- In some locations, sprinkler irrigation is used for frost or freeze protection, or for crop cooling. In these special cases, applications should be limited to the amount necessary for crop protection, and applied water should remain on site.

Appendix C. Enforcement Procedures

The Oregon Department of Agriculture's primary mission is to support Oregon's agricultural industry. The Oregon Department of Agriculture will have the responsibility for enforcing rules derived from the Curry County Agricultural Water Quality Management Area Plan. It is the intent of the Local Advisory Committee and ODA policy that fines and civil penalties be used as a last resort in the effort to improve water quality in Curry County. This is consistent with the direction given to the Department through the Oregon Administrative Rules for the Agricultural Water Quality Management Program (603-090-0000 through 603-090-0120). This Area Plan includes an enforcement policy because it is a required element of a Water Quality Plan and to provide a mechanism when reasonable attempts at voluntary solutions have failed.

The primary focus of the Curry County AgWQM Area Plan is education toward voluntary compliance with the plan. Even the enforcement procedure is designed to educate first and penalize only as a last resort.

In the event that a situation comes to the attention of the Oregon Department of Agriculture that may be a violation of the Curry County Agricultural Water Quality Administrative Rules, a prescribed procedure will be followed. EXCEPT FOR BLATANT POLLUTION OF WATERS OF THE STATE OR BLATANT DESTRUCTION OF ADEQUATE RIPARIAN VEGETATION ALONG PERENNIAL STREAMS, AT ANY POINT IN THE PROCESS, THE LANDOWNER MAY CHOOSE TO ADDRESS A PROBLEM AND NO CIVIL PENALTIES WILL BE LEVIED BY THE OREGON DEPARTMENT OF AGRICULTURE.


1. Any person alleging a violation of the Curry County Agricultural water quality administrative rules may file a complaint with the Oregon Department of Agriculture. The department will evaluate or investigate a complaint filed by a person if the complaint is in writing, signed, and dated by the complainant, and indicates the location and description of the violation of the Curry County Agricultural Water Quality Administrative Rules.
2. If the problem appears to be a violation of the Curry County Agricultural Water Quality Administrative Rules an Oregon Department of Agriculture representative will contact the landowner to schedule a meeting. NO OREGON DEPARTMENT OF AGRICULTURE REPRESENTATIVE WILL ENTER PRIVATE PROPERTY AT ANY TIME WITHOUT THE OWNER'S PERMISSION OR A VALID SEARCH WARRANT.
3. The situation will be reviewed on-site by an Oregon Department of Agriculture representative and the landowner. The on-site review will include an investigation by the Oregon Department of Agriculture, which may include collecting appropriate samples for testing and consultation with experts as appropriate, at the Oregon Department of Agriculture's expense. If no problem exists, the complaint would be dropped.
4. If the Oregon Department of Agriculture determines through the investigation that a violation of the Curry County Agricultural Water Quality Administrative Rules exists, the Oregon Department of Agriculture will advise the landowner of the violation (i.e. issue a notice of noncompliance) and work with the landowner to develop a plan of correction to solve the problem. The plan of correction includes a timetable and an agreement to revisit the site as necessary to confirm that progress is being made to correct the violation within the timetable agreed upon. This would complete the process.

5. If the landowner does not agree that a problem exists, the landowner may choose to do additional testing or consultation at their own expense and request a review by the department of the initial findings in light of any additional information collected.
6. If evaluation of the additional information by the Oregon Department of Agriculture determines that no problem exists or that the violation is not the result of an agricultural practice by the landowner, the process is complete.
7. If there is a confirmed problem that a landowner refuses to address after the department's on-site visit and the department's attempts to work with the landowner to develop a mutually agreeable solution, civil penalties can be levied. Civil penalties are issued by the Oregon Department of Agriculture Director or the director's designee and will be based on the seriousness of the violation and the magnitude of the effect. OAR 603-090-0120(3) describes the civil penalty matrix for first violations which begins at \$50.00 and ranges to \$1200.00, and the civil penalty matrix for repeat violations which begins at \$100 and ranges to \$5000.00. ORS 568.933 states 'each day of violation continuing after the period of time for correction set by the department shall be considered a separate violation unless the department finds that a different period of time is more appropriate to describe a specific violation event.'
8. A landowner issued a civil penalty due to a violation of the Curry County Agricultural Water Quality Administrative Rules may request a hearing with the Director of the Department of Agriculture. The hearing provides for the director to hear the landowner's disposition from which the director determines appropriate action, which can include a modification of the civil penalty or other form of intermediate sanction.
9. A landowner issued a civil penalty due to a violation of the Curry County Agricultural Water Quality Administrative Rules may request a formal hearing by a hearings officer assigned from the Hearing Officer's Panel in accordance with applicable contested case procedures as described in ORS 183.413 to 183.550. Upon conclusion of the hearings process, a hearings officer will prepare a proposed order that includes recommended findings of fact, conclusions of law, and appropriate action by the agency. If the order is in favor of the landowner, the process is complete. If not, the landowner becomes subject to procedures for payment of the civil penalty.

NO CIVIL PENALTIES WILL BE ISSUED DURING THE FIRST TWO YEARS FOLLOWING PLAN ADOPTION, UNLESS THERE IS BLATANT POLLUTION OF WATERS OF THE STATE OR BLATANT DESTRUCTION OF ADEQUATE RIPARIAN VEGETATION ALONG PERENNIAL STREAMS.

THE PROCESS IS DESIGNED TO BE FAIR TO THE LANDOWNER AND TO ALLOW ENFORCEMENT OF THE CURRY COUNTY AGRICULTURAL WATER QUALITY ADMINISTRATIVE RULES.

ODA FORMAL WATER QUALITY COMPLAINT PROCEDURE



*As determined by scientific means

*As determined by Scientific Means

Appendix D. Map of Curry County Agricultural Water Quality Management Area

